

# FOXP3(JF0926)

rev. 01/09/17  
Cat#: ET1702-16

**Product Type:** Recombinant rabbit monoclonal IgG, primary antibodies

**Species reactivity:** Human

**Applications:** WB, FC

**Molecular Wt.:** 50 kDa

**Clone number:** JF0926

**Description:** The FOX family of transcription factors is a large group of proteins that share a common DNA binding domain termed a winged-helix or forkhead domain. During early development, FOXP1 and FOXP2 are expressed abundantly in the lung, with lower levels of expression in neural, intestinal and cardiovascular tissues, where they act as transcription repressors. FOXP1 is widely expressed in adult tissues, while neoplastic cells often exhibit a dramatic change in expression level or localization of FOXP1. The gene encoding human FOXP1 maps to chromosome 3p14.1, and the gene encoding human FOXP2 maps to chromosome 7q31. The gene encoding FOXP3, a third member of this family, maps to chromosome Xp11.23. Mutations in this gene cause IPEX, a fatal, X-linked inherited disorder characterized by immune dysregulation. The FOXP3 protein, also known as scurf, is essential for normal immune homeostasis. Specifically, FOXP3 represses transcription through a DNA binding forkhead domain, thereby regulating T cell activation.

**Immunogen:**

Recombinant protein.

**Positive control:**

Raji.

**Subcellular location:**

Nucleus, Cytoplasm.

**Database links:**

SwissProt: Q9BZS1 (Human)

**Recommended Dilutions:**

**WB:** 1:500-1:1,000

**FC:** 1:10-1:50

**Storage Buffer:**


1\*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.

**Storage Instruction:**

Store at +4° C after thawing. Aliquot store at -20° C or -80° C. Avoid repeated freeze / thaw cycles.

**Purity:**

ProA affinity purified.



**Fig1:** Flow cytometric analysis of Raji cells with FOXP3 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

**Background References**

1. Nguyen N et al. Tumor infiltrating lymphocytes and survival in patients with head and neck squamous cell carcinoma. *Head Neck* 38:1074-84 (2016).
2. Gu L et al. Rapamycin ameliorates CCl4-induced liver fibrosis in mice through reciprocal regulation of the Th17/Treg cell balance. *Mol Med Rep* 14:1153-61 (2016).

Hangzhou HuaAn Biotechnology Co.,Ltd.

Orders: 0086-571-88062880

Support: 0086-571-89986345

Service mail: tech@huabio.com

www.huabio.com



华安生物

Applications: WB=Western blot IP=Immunoprecipitation IHC=Immunohistochemistry IF=Immunofluorescence FC=Flow cytometry  
Species Cross-Reactivity: H=human M=mouse R=rat Hm=hamster Mk=monkey Mi=mink C=chicken Dm=D.melanogaster X=Xenopus Z=zebrafish  
B=bovine Dg=dog Pg=pig Sc=S.